

TABLE 1. MICs of 20 antimicrobial agents tested for five strains of *A. baumannii*

| Strain | MIC range ($\mu\text{g/ml}$) ^a | | | | | | | | | | | | | | | | | | | |
|----------|---|-----|-------|-----|------|-----|-----|------|------|-------|------------|-----|-----|------|-------|-----|---------|-----|-----|-----|
| | GEN | TOB | NET | AMK | NAL | CIP | OFX | AMP | PIP | SAM | AMC | CXM | CTX | CAZ | IPM | MEM | SXT | TET | CHL | COL |
| RUH 875 | >32 | >16 | 32 | 8 | 16 | 1 | 1 | >128 | 256 | 32/16 | 16/8-32/16 | 64 | 32 | 8 | 0.5-1 | 1 | >8/152 | >64 | >64 | 0.5 |
| RUH 134 | 32 | 2 | 8 | 2-4 | 8 | 0.5 | 0.5 | >128 | 256 | 32/16 | 16/8 | 32 | 16 | 8 | 0.5 | 0.5 | >8/152 | >64 | >64 | 0.5 |
| LUH 6034 | >32 | 8 | >64 | 64 | >128 | >32 | 16 | >128 | >256 | 64/32 | >128/64 | >64 | >64 | >128 | >32 | >32 | 8/152 | >64 | >64 | 0.5 |
| LUH 7312 | >32 | 4 | 16-32 | 64 | >128 | >32 | 16 | >128 | 64 | 8/4 | 16/8 | 64 | 32 | 8 | 0.5-1 | 1 | >8/152 | >64 | >64 | 0.5 |
| LUH 7858 | 2-4 | 1 | 8-16 | 128 | >128 | >32 | 32 | >128 | >256 | 16/8 | >128/64 | >64 | >64 | >128 | >32 | 16 | 0.5/9.5 | 64 | >64 | 0.5 |

^a Abbreviations: GEN, gentamicin; TOB, tobramycin; NET, netilmicin; AMK, amikacin; NAL, nalidixic acid; CIP, ciprofloxacin; OFX, ofloxacin; AMP, ampicillin; PIP, piperacillin; SAM, ampicillin-sulbactam; AMC, amoxicillin-clavulanic acid; CXM, cefuroxime; CTX, cefotaxime; CAZ, ceftazidime; IPM, imipenem; MEM, meropenem; SXT, trimethoprim-sulfamethoxazole; TET, tetracycline; CHL, chloramphenicol; COL, colistin.